

# Osmotic performance rate, stress response and growth performance of silver pompano (*Trachinotus blochii*) reared in different salinities using recirculating culture system

<sup>1</sup>Mulyadi, <sup>1</sup>Usman M. Tang, <sup>2</sup>Bintal Amin, <sup>1</sup>Sukendi, <sup>1</sup>Niken A. Pamukas, <sup>3</sup>Windarti

<sup>1</sup> Department of Aquaculture, Faculty of Fisheries and Marine Science, Riau University, Pekanbaru, Indonesia; <sup>2</sup> Department of Marine Science, Faculty of Fisheries and Marine Science, Riau University, Pekanbaru, Indonesia; <sup>3</sup> Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Riau University, Pekanbaru, Indonesia. Corresponding author: Mulyadi, mulyadibrian26@yahoo.com

**Abstract.** Silver pompano (*Trachinotus blochii*) has received tremendous attention from the aquaculture sector, due to its favourable features, such as a high economic value, its good adaptive response and its potential to be cultured in various salinities. The aim of this study was to discover the effects of medium salinities on osmotic performance rate, blood cortisol, and growth performance (absolute growth weight and length, specific growth rate (SGR), feed conversion (FC), feed efficiency (FE), survival rate (SR)) of *T. blochii* under recirculating system. The histological alterations (kidney and gill) and water quality (temperature, pH, DO, NH<sub>3</sub>, NO<sub>2</sub> and NO<sub>3</sub>) were also observed. The 56-day experiment was carried out in Balai Perikanan Budidaya Laut (BPBL) of Batam, Indonesia. A total of 300 fish specimens (11-13 cm in length, weighing 28-29 g) were raised in a 100 L tank containing 80 L of water at a density of 1 fish 4 L<sup>-1</sup> (20 fish in total). They were fed with commercial pellet (46% protein) at 3% of fish biomass and 3 times a day. The experiment was conducted according to a completely randomized design with 5 levels of salinity: P<sub>1</sub>=25‰, P<sub>2</sub>=20‰, P<sub>3</sub>=15‰, P<sub>4</sub>=10‰ and P<sub>5</sub>=5‰, by performing triplicate measurements for each treatment. The treatment with 15 ‰ salinity showed the best effects, yielding an osmotic performance rate of 3 mOsm L<sup>-1</sup> H<sub>2</sub>O, a blood cortisol level of 50,923 nmol L<sup>-1</sup>, an absolute growth weight and length of 17.73±1.25 g and 2.32±0.21 cm, respectively, an SGR of 0.87±0.05%, an FC of 1.24±0.00, an FE of 80.79±0.58 and an SR of 88.33±2.88%. Histologically, there were no anomalies in the structure of gill and kidney of the fish cultured in 5‰-25‰ salinities. Water quality was acceptable for growing *T. blochii*.

**Key Words:** gill, kidney, blood cortisol, histology, water quality.

**Introduction.** The farming of silver pompano (*Trachinotus blochii*) has currently gained great popularity in Indonesia. The market demand for the species has continuously increased in international trade, due to its high economic value, its good adaptive response and its potential to be cultured in various water conditions. The price of the fish reaches approximately USD 4.25 kg<sup>-1</sup> in local market, but it may reach USD. 14.144 kg<sup>-1</sup> in export market (Mo 2017). Since 2015, *T. blochii* is considered as a promising commodity in marine fisheries sector with total production of 1,900 tons in 2015, demonstrating the annual rise of 31.5% (Prahadi 2015).

*T. blochii* was reported to exert a high adaptive response towards salinity changes. Survival rate of the species at various salinity levels 32‰, 24‰, 14‰, and 4‰ during the 28-day trial showed no significant difference, indicating that the fish could adapt at lower salinity over seawater, as well as confirming the possibility of fish farming diversification through culture systems in brackish water (Arrokhman et al 2012).

Salinity tolerance in fish closely relates to the osmotic pressure balance between inside and outside the fish body. Osmotic pressure inside the fish is lower than outside.

The imbalance may cause disturbance of the physiological functions, which in turn disrupt the fish growth. However, under a normal osmotic gradient, the metabolic activity could reach an optimum rate, as indicated by a good appetite and an enhanced feed intake, allowing allocating more energy for the growth (Carrion et al 2005). Fujaya (2004) suggested that the osmotic balance was achieved through regulation of body fluid transportation, known as osmoregulation. The adjustment activities needed for balancing internal and external osmotic pressures require a high energy consumption, leading to stress generation, as indicated by the production of blood cortisol.

Important osmoregulatory organs including gills and kidney play a crucial role in the process. Defective tissues in these organs offer a sign of failure in osmoregulation. Consequently, the defect would adversely affect physiological functions, causing the decrease of feed consumption and fish growth (Putri et al 2014). In this matter, gills seem to be the most susceptible osmoregulatory organ towards environmental changes, such as physicochemical properties of water and presence of toxic compounds. The gill lamellae become the weakest part, in which presence of stressors directly induces ionic homeostasis that remarkably imparts osmoregulation. Indeed, the chronic stressors lead to destructive effects on the gill. Macroscopic and microscopic defects in the gill can serve as biomarker of fish health status (Camargo & Martinez 2007). Besides, Thophon et al (2003) also argued that kidney is a susceptible organ to the external stressors exposure, due to its essential role in maintaining homeostasis. Based on the above elaboration, it can be concluded that there is a need for investigating the osmotic response, stress level and growth performance of *T. blochii* reared in a recirculation system, at different salinity levels. In this work, the histological alterations in gill and kidney were also observed.

## Material and Method

**Study site.** The experiment was carried out in the Agency for Marine Fisheries Culture (Balai Perikanan Budidaya Laut-BPPL), Batam, Indonesia, for 56 days. A total of 300 *T. blochii* seeds (average length of 11-13 cm, weight of 28-29 g) were reared in an experimental container (capacity 100 L) filled with 80 L of water. The density referred to Indonesia National Standard (SNI 2013), which recommends 1 individual 4 L<sup>-1</sup> (equal to 20 individuals 80 L<sup>-1</sup>). Commercial pellet (Megami GR 2) was used, containing 46% protein, 9% fat, 1.9% crude fiber and 8% moisture. The specimens were fed at 3% of their weight, three times a day.

**Preparation of the culture container.** The close recirculating system was prepared. The container was filled with seawater at different salinity levels, then connected to PVC gutter (50 x 14 x 14 cm) at the upside of the chamber. The filtration unit water was transported into the culture container through a PVC pipe (2.5 cm diameter). The filtration unit was filled with 50 bioballs (each gutter), as previously prescribed by Nelvia et al (2015). The water was subsequently pumped to the filtration unit with the aid of a 50 W water pump.

**Experimental design.** The completely randomized design was arranged, consisting of 1 factor and 5 levels (with triplicates), as follows: P<sub>1</sub>=25‰, P<sub>2</sub>=20‰, P<sub>3</sub>=15‰, P<sub>4</sub>=10‰ and P<sub>5</sub>=5‰. The effect of salinity was studied, focusing on the osmotic response, the content of blood cortisol, the tissue histology (gills and kidney), the absolute growth weight (Wm), the growth length (Lm), the specific growth rate (SGR), the survival rate (SR) and the water quality (temperature, pH, DO, NH<sub>3</sub>, NO<sub>2</sub> and NO<sub>3</sub>).

**Determination of the osmotic response.** Osmotic response was evaluated at the end of experiment and determined using micro-osmometer, as explained by Cambell et al (2012). The fish sample was acclimatized for 7 days and, for the first 5 days, the fish was fed. A total of 18 fish specimens were used for each salinity level, in which they were exposed to the salinity treatments. After 7 days of exposure, the blood of living fish (1 mL) was collected from the heart, and then centrifuged at 3,000 rpm for 3 minutes to obtain blood plasma. The plasma osmotic pressure was measured using a micro-osmometer. The osmoregulatory capacity refers to the difference between osmotic

pressure of the medium and internal osmotic pressure of the fish blood plasma. The calculation was described as follows (Anggoro & Nakamura 1996):

$$\text{TKO} = [\text{POsmoBlood} - \text{POsmoMedium}]$$

Where:

TKO - osmotic response ( $\text{mOsm L}^{-1} \text{H}_2\text{O}$ );

POsmoBlood - biota osmolarity ( $\text{mOsm L}^{-1} \text{H}_2\text{O}$ );

POsmoMedium - medium osmolarity, and the bracket [ ] means absolute value.

**Blood cortisol analysis.** The analysis was performed at 3 periods: day 1, day 28 and day 56 of the experiment, using enzyme-linked immunosorbent assay. Before the blood withdrawal, the fish was anaesthetized using phenoxyethanol at a dose of  $0.3 \text{ mL L}^{-1}$  water (Rigal et al 2008). Briefly, blood was collected via the vena caudalis, using a heparinized syringe (1 mL), then centrifuged to collect plasma. Plasma cortisol was quantified using RIA (radio immunoassay) Cortisol ( $^{125}\text{I}$ ) RIA KIT IZOTOP (Ramsay et al 2006). In this regard, blood plasma (0.3 mL) was frozen at  $-20^\circ\text{C}$ . To maintain the hormone in the plasma, the sample was packed within a cool-box containing dry ice exactly at day 57 of experiment, then immediately transported into laboratory for analysis.

**Histological analysis.** Gill and kidney tissue of the sample was collected at the first and last days of the experiment, from the fish specimens exposed to a salinity of 30 ppt (natural habitat condition). Gill cover (overculum) was lifted and the base was cut off to release the gill. Afterwards, the kidney was obtained by opening the abdominal cavity. The phosphate-buffered formalin (NBF) at 10% was used for organ fixation, carried out for 24-48 h (Raškoviæ et al 2011). After fixation, the tissues were dehydrated in graded series of alcohol and xylol, and then embedded in paraffin. All these stages were conducted using tissue processor. The sections obtained were cut at thickness of 3-5  $\mu\text{m}$ , then incubated in a water bath at  $40^\circ\text{C}$  and then immediately air-dried for 1 h. Afterwards, it was stained using haematoxylin-eosin (HE), and observed under light microscope at magnification of 400 $\times$ .

**Growth performance.** The weight and length of fish were recorded each 14 days. Growth performance included following parameters:

- a. Weight gain (g) = final weight - initial weight;
- b. Absolute length growth (cm) = average final length (cm) - average initial length (cm);
- c. Specific growth rate (SGR) (%) =  $(\text{Ln mean final fish weight} - \text{Ln mean initial fish weight}) / \text{culture period (day)} \times 100$ ;
- d. Feed efficiency (FE) (%) = increased fish mass/total feed consumed;
- e. Survival rate (%) = (final number of fish/initial number of fish)  $\times 100$ .

**Water quality.** Temperature was observed daily using a thermometer, while chemical indicators were checked each 14 days, including pH (using pH meter), DO (using DO meter),  $\text{NH}_3$ ,  $\text{NO}_2$  and  $\text{NO}_3$  (using a spectrophotometer).

## Results and Discussion

**Osmotic performance rate.** Osmotic response of fish to the variation of salinity was shown in Figure 1, indicating that salinity caused a remarkable effect on medium osmolarity, blood plasma, and osmotic performance rate of *T. blochii* ( $p < 0.05$ ). In this case, the lowest rate ( $3 \text{ mOsm L}^{-1} \text{H}_2\text{O}$ ) was found at P3 (15‰), while the highest one ( $87 \text{ mOsm L}^{-1} \text{H}_2\text{O}$ ) was attributed to P5 (5‰). This clearly imparts the effect of salinity on medium and plasma osmolarity, leading to noticeable changes in osmotic performance rate of *T. blochii* seeds. The extreme difference between medium osmolarity and internal fish osmolarity could cause significant changes in fish behaviors and physiological conditions, consequently altering the feed consumption rate and fish growth.

In water of salinity near the isosmotic condition (15‰), the fish allocated more energy for enhancing their growth. Energy metabolism required for osmoregulation in fish is fundamentally associated with osmotic performance rate as a rapid response towards changes in medium osmolarity. In this case, osmotic performance rate shows linear correlation to energy consumption for osmoregulatory activities. On the contrary, hypoosmotic (salinity of 30, 25 and 20‰) and hyperosmotic (10 and 5‰) condition leads to increment of osmotic response, which needs higher energy requirement for osmoregulation (Carrion et al 2005). Arjona et al (2009) reported that there was clear evidence that higher osmotic response led to higher energy use for osmoregulation. Our data demonstrated a variety of osmotic response with progression of salinity. As mentioned by Putri et al (2014), medium that possesses higher salinity (far away from isosmotic condition) would increase osmotic performance rate of the fish. At the condition in which the medium condition is tolerable, osmotic performance rate tended to attenuate, thus allowing energy use for fish growth.




Figure 1. Average osmolarity of rearing medium and blood plasma, and osmotic performance rate of *Trachinotus blochii* cultured in various salinity levels. Different superscripts above the bar showed significant difference at  $p < 0.05$ .

This present work reveals that *T. blochii* is able to adapt to the medium salinity at 15 ppt, suggesting that the juvenile (average length of 12.43-12.48 cm, weight 28.00-28.70 g and age of 2 months) can tolerate the condition. Additionally, Retnani & Abdulgani (2013) argued that such adaptive capability of fish relied on size and growth stage, in which osmoregulatory activity of fish may differ depending on age. Lantu (2010) stated that osmoregulation of euryhaline fish strongly related to osmo-sensitivity of chloride cells. They serve as receptor, mainly responding to the salinity level of the medium. When immersed in water media with different salinity, chloride cells in euryhaline fish transmit signals to central nervous system, primarily to the pituitary gland responsible for controlling secretion of growth hormone. Subsequently, the hormone regulates the development of chloride cells in osmoregulatory organs such as gills, kidney and digestive tract. Thus, the amount of chloride cells was adjusted, causing changes in the physiological mechanisms of secretion or absorption of ions by chloride cells. At a higher salinity, chloride cells would have a higher rate of proliferation, and vice-versa. Over the long term, this controlling mechanism may also cause genetic expression.

Syakirin et al (2018) stated that a higher concentration of ions in water would rise salinity level and osmolar density. For instance, the hybrid grouper is a kind of marine fish that has blood osmolarity (internal osmotic fluid pressure) lower than the environmental osmotic pressure. Therefore, the water will pass from the body of the fish to the environment by the osmotic process through the kidney, gill, as well as in the body. Salinity demonstrates a relationship with the osmoregulation of aquatic animals. A sudden fluctuation of salinity makes the body osmoregulation difficult and induces the animal mortality. The osmoregulation capacity is determined by the difference between the blood osmotic pressure (fish) and the media osmotic pressure. Osmoregulation relates to the difference of fish blood osmolarity and media osmolarity, known as the osmotic work level: the response to the salinity change increases with this difference.

**Stress level in *T. blochii*.** The increment of blood cortisol in fish indicates stressful condition. Our data suggested that blood cortisol in all treated samples declined at different extent during the experiment progression, indicating that the fish exhibited adaptive capacities to lower salinity (Figure 2).




Figure 2. Concentration of blood cortisol in *Trachinotus blochii* cultured in various levels of salinity.

As depicted in Figure 2, the concentration of blood cortisol on day 1 increased as the level of salinity was lower compared to the control (30‰). This clearly showed a stress response to the new environment. Setiyoningsih (2014) argued that stress in fish existed as rapid response towards environmental pressure, thus they secreted glucocorticoid (cortisol) and catecholamine hormone to cope with the stress condition.

Furthermore, cortisol level declined consistently from day 28 to day 56 in all treatments, indicating adaptive capacities. On day 56, the lowest cortisol level (50,923 nmol L<sup>-1</sup>) was found at 15‰, while the highest one (94,160 nmol L<sup>-1</sup>) was found at 5‰. In salinity 15‰, the fish showed the best osmoregulatory activity through balancing osmotic pressures. Pamungkas (2012) argued that osmoregulation in fish was modulated by two hormones, namely prolactin and cortisol. Cortisol is a crucial hormone in euryhaline fish since it modulates excretion of ions via gills able to stimulate chloride cells, thus, when migrating, the concentration of plasma cortisol increases. In addition,

Scabra (2018) found the depletion of blood volume, leukocyte, and liver glycogen in stressed fish, but the concentration of cortisol increased. Stress condition results mainly from external changes such as salinity, and during this stressful period, fish activate homeostatic processes by accelerating their metabolic activities, leading to the rise of oxygen intake.

At the end of experiment, the concentration of cortisol in 15, 20 and 25‰ salinity reached a level close to the value measured for the control salinity (30‰). This represents the successful attempt of the fish to reach stability. Hastuti et al (2004) reported that concentrations of cortisol in plasma of normal fish ranged between 20.65–53.22 µg dL<sup>-1</sup>.

**Water quality.** Table 1 presents the parameters of water quality, including temperature, dissolved oxygen (DO), pH, ammonia (NH<sub>3</sub>), nitrite (NO<sub>2</sub>), and nitrate (NO<sub>3</sub>). The results showed that these parameters tended to be similar in all salinity levels, within the following ranges: temperature 27-29°C, pH 5.9-7.9, DO 5.9-10.9 mg L<sup>-1</sup>, NH<sub>3</sub> 0.01–0.131 mg L<sup>-1</sup>, NO<sub>2</sub> 0.050–0.090 mg L<sup>-1</sup>, and NO<sub>3</sub> 0.190–0.890 mg L<sup>-1</sup>. It is noticeable that these values correspond to a set of good conditions for growth and survival rate of *T. blochii*. As discussed by Sitta & Hermawan (2011), the optimum condition for *T. blochii* included temperature 28-32°C, pH 6.8-8.4, while DO ranged 4.8–5 mg L<sup>-1</sup> in aquarium and ±7.3 mg L<sup>-1</sup> in floating net cage. Boyd (2015) found that ammonia at level of 0.2–2.0 mg L<sup>-1</sup> could be detrimental to fish, while nitrite was acceptable for fish at a concentration <1 mg L<sup>-1</sup>, unsafe at 1-5 mg L<sup>-1</sup>, and poisonous at 16 mg L<sup>-1</sup> (Siikavuopio & Saether 2006).

Table 1

Mean water quality parameters during the research period

Parameters	Unit	Salinity				
		25‰	20‰	15‰	10‰	5‰
Temperature	°C	27.3-28.8	27.4-28.6	27.8-29.1	27.3-28.6	27.5-28.6
pH	-	7.4-7.9	7.0-7.9	6.9-7.9	6.1-7.9	5.9-7.8
DO	mg L <sup>-1</sup>	5.9-6.6	8.8-10.9	5.9-6.8	5.9-7.03	6-7.1
NH <sub>3</sub>	mg L <sup>-1</sup>	<0.01- 0.105	<0.01- 0.101	<0.01- 0.099	0.010- 0.109	0.060- 0.131
NO <sub>2</sub>	mg L <sup>-1</sup>	0.050- 0.070	0.050- 0.059	0.050- 0.059	0.060- 0.071	0.060- 0.090
NO <sub>3</sub>	mg L <sup>-1</sup>	0.340- 0.750	0.360- 0.780	0.360- 0.890	0.190- 0.570	0.206- 0.345

During the experiment, water quality was maintained at a large extent to ensure acceptable parameters for fish growth. Water filtration was installed in the culture system, comprising of physical (synthetic cotton), chemical (zeolite and active carbon), and biological (bioball) filter. Cotton filter served to capture uneaten feed and feces, while zeolite and active carbon enabled the absorption of toxic compounds such as ammonia and nitrite (Supriyono et al 2007). Bioball is important as attachment site for nitrifying bacteria capable of converting nitrogen into unharmed form, i.e. nitrate (Dewi & Masithoh 2013). Nurhidayat et al (2012) also augmented that the combination of zeolite, active carbon, and bioball showed satisfying results of maintaining water quality through oxidation of ammonia and enrichment of non-pathogenic nitrifying bacteria.

**Growth performance and survival rate.** Salinity demonstrated significant impacts to absolute growth weight, absolute growth length, SGR, feed conversion, FE, and SR of *T. blochii* (p<0.05). Statistical test of Newman-Keuls revealed that two salinity levels, i.e. 5‰ and 10‰, did not result in any significant difference in some parameters including absolute growth weight and growth length, SGR, feed conversion, and FE. Meanwhile, SR tended to be similar between treatments (Table 2).

Table 2

Growth performance of *Trachinotus blochii* reared in different salinity levels

Parameters	Salinity				
	25‰	20‰	15‰	10‰	5‰
Absolute weight growth (g)	14.03±1.18 <sup>ab</sup>	15.87±1.05 <sup>bc</sup>	17.73±1.25 <sup>c</sup>	12.73±1.70 <sup>a</sup>	11.93±1.66 <sup>a</sup>
Absolute length growth (cm)	1.93±0.34	1.82±0.14	2.32±0.21	2.09±0.23	1.84±0.07
Specific growth rate (%)	0.75±0.08 <sup>ab</sup>	0.87±0.04 <sup>b</sup>	0.87±0.05 <sup>b</sup>	0.67±0.09 <sup>a</sup>	0.62±0.07 <sup>a</sup>
Feed conversion	1.31±0.01 <sup>a</sup>	1.26±0.01 <sup>a</sup>	1.24±0.00 <sup>b</sup>	1.33±0.01 <sup>c</sup>	1.34±0.01 <sup>d</sup>
Feed efficiency (%)	76.62±0.8 <sup>a</sup>	79.07±1.03 <sup>a</sup>	80.79±0.58 <sup>b</sup>	75.22±0.76 <sup>c</sup>	74.37±0.28 <sup>a</sup>
Survival rate (%)	81.67±2.88 <sup>ab</sup>	83.33±2.88 <sup>ab</sup>	88.33±2.88 <sup>b</sup>	83.33±2.88 <sup>ab</sup>	76.67±2.88 <sup>a</sup>

a, b, c and d were significantly different; ab and bc were not significantly different ( $p < 0.05$ ).

As presented in Table 2, 15‰ salinity demonstrated the most satisfying effects on growth of *T. blochii*, meaning that the fish could perform proper feed utilization and osmoregulation. Wulandari (2006) reported that optimum energy use could be achieved at osmotic condition; thus, more energy was used for their growth instead of osmoregulatory activities.

Retnani & Abdulgani (2013) reported that growth of *T. blochii* cultured in 4-24‰ salinity was better than that in 32-34‰ salinity. The salinity modification to less than the seawater salinity level provoked a decline of the used energy owing to the attenuation of ion exchanges by gill's chloride cells. Such a condition minimized the energy demand for osmoregulation, thereby enhancing the fish growth.

Considering that size, age, stock density and feed are similar, the difference of the fish growth is undoubtedly affected by the environmental salinities. We also noted that water quality ranged within value intervals appropriate for the culture of *T. blochii*. The importance of salinity for the fish growth is associated with changes in physiological functions. Fish cultured in high salinity (control) performed a more active transport in order to release excessive ions of Na from gill, which is a highly energy-consuming activity. Gill chloride cells are responsible for fish osmoregulation. Proliferated on the lamellae, they are extremely sensitive to external salinity. When moving to the new medium with different salinity, euryhaline fish activated chloride cells and delivered signals to central nervous system. In a culture medium with higher salinity, the proliferation of chloride cells was more intensive; conversely, they were less produced under lower salinity conditions (Bone & Moore 2008; Fujaya 2004).

The specific growth rate (SGR) was the highest in fish cultured in 15‰ and 20‰ salinity (0.87%), and the lowest in 25‰ (0.75%), 10‰ (0.67%) and 5‰ (0.62%) salinity, respectively. Retnani & Abdulgani (2013) reported a higher SGR in *T. blochii* cultured in 24‰ salinity (10.594%) than in 32-34‰ salinity (10.359%). Noticeably, the SGR of *T. blochii* farmed in brackish salinity ranged from 14‰ to 24‰.

Feed efficiency (FE) reached the highest level in fish cultured at 15‰ salinity, indicating that isosmotic condition was achieved. Therefore, energy expenditure is devoted to fish growth instead of osmoregulation. Febrianti et al (2016) argued the efficient utilization of feed must be greater than 50%. We further noted that FE of *T. blochii* exceeded 50% across the treatments, meaning that the fish could utilize the feed efficiently in various salinities.

In respect of the survival rate (SR), the percentage recorded across the treatments ranged within 76.67-88.33%, while fish cultured at 15‰ salinity reached the highest level. It is noticeable that *T. blochii* can properly adapt to a wide range of salinities, i.e. 30‰-5‰. Additionally, Arrokhman et al (2012) reported a SR of 99.03–100% in *T. blochii* reared in 4-34‰ salinity. This suggests that the fish has the potential to be farmed in brackish water.

**Histological alterations.** Histological observation on the fish gills did not find anomalies in all treatments studied, as indicated by the clear appearance of the secondary lamellae, epithelium, thrombocytes and pillar cells. No abnormalities were observed in gill chloride cells, including the absence of hypertrophy and edema, despite a moderate hyperplasia (Figure 3). In this case, hyperplasia was linked to parasites in the medium. As reported by Wahyuni et al (2017), normal tissue of fish gill was characterized by obvious appearance of secondary lamellae, pillar cells, lacunae, and thrombocytes.




Figure 3. Structure of gill tissue of *Trachinotus blochii* cultured in different levels of salinity. K-control, P<sub>1</sub>=25‰, P<sub>2</sub>=20‰, P<sub>3</sub>=15‰, P<sub>4</sub>=10‰ and P<sub>5</sub>=5‰, sp-pillar cells, ls-secondary lamellae, ep-epithelium, sd-thrombocyte, hp-hyperplasia.

The histological analysis on fish kidney did not find any significant abnormalities, as indicated by condition of Bowman's capsule and glomerulus. The cells were intact and not infected, but a bleeding part was observed (Figure 4). Mc Gavin & Zachary (2007) offered a description of the kidney histology comprised of main parts such as glomerulus, tubulus and blood vessels. In addition, Takashima & Hibiya (1995) reported that Bowman's capsule surrounds glomerulus as an indicator of healthy kidney. These organs perform an essential role, i.e. filtering metabolites in bloods. The excretory fluids enter the tubule, while minerals, glucose, and other fluids are re-absorbed. The number and size of glomerulus in freshwater fish were greater than those in seawater fish, considering their importance in retaining salt in body and releasing urine.






Figure 4. Structure of kidney tissue of *Trachinotus blochii* cultured in different levels of salinity. K-control, P<sub>1</sub>=25‰, P<sub>2</sub>=20‰, P<sub>3</sub>=15‰, P<sub>4</sub>=10‰ and P<sub>5</sub>=5‰, Kb-Bowman's capsule, G-glomerulus, P-bleeding.

**Conclusions.** The experimental data revealed the significant effects of salinity on the osmoregulatory activities, blood cortisol level, and growth performance of *T. blochii* ( $p < 0.05$ ). The variance of salinities did not cause any difference in the structure of gill and kidney and did not show any abnormality in these organs. The treatment at 15‰ salinity exhibited the best outcome, resulting in: osmotic pressure of 3 mOsm L<sup>-1</sup> H<sub>2</sub>O (closer to the isosmotic condition), blood cortisol of 50,923 nmol L<sup>-1</sup>, absolute growth weight of 17.73±1.25 g, absolute growth length of 2.32±0.21 cm, specific growth rate of 0.87±0.05%, feed conversion of 1.24±0.00, feed efficiency of 80.79±0.58% and survival rate of 88.33±2.88%. Parameters of water quality were at the appropriate level for *T. blochii* growing. Furthermore, the fish could exert adaptive capacities to medium salinities below the seawater salinity, the most satisfying level being achieved at 15‰ salinity, when cultured in a recirculation system.

**Acknowledgements.** Authors would like to acknowledge the Agency for Marine Fisheries Culture, Batam for providing research facilities. We would also like to thank the Riau University Research Centre, Riau University and the Ministry of Research, Technology and Higher Education for research funding, as well as Razuli, Eva Selvia, Hidayati and Fitriulan for technical assistance.

## References

- Anggoro S., Nakamura K., 1996 Osmoregulation of Kuruma prawn (*Penaeus japonica*). Bulletin Kagoshima 2(3):14-19.
- Arjona J. F., Chacoff L. V., Jarabo I. R., Gonçalves O., Páscoa I., María P., Río M. D., Mancera J. M., 2009 Tertiary stress responses in Senegalese sole (*Solea senegalensis* Kamp. 1858) to osmotic challenge: implication for osmoregulation, energy metabolism and growth. Aquaculture 287:419-426.
- Arrokhman S., Abdulgani N., Hidayati D., 2012 [Survival rate of silver pompano (*Trachinotus blochii*) reared in culture media using different salinity]. Jurnal Sains dan Seni ITS 1(1):32-35. [In Indonesian].
- Boyd C. E., 2015 Water quality. An introduction. Springer International Publishing, AL, USA, 374 p.

- Bone Q., Moore R. H., 2008 Biology of fishes. Taylor & Francis Group 161-171:260-271.
- Camargo M. M. P., Martinez C. B. R., 2007 Histopathology of gills, kidney and liver of a neotropical fish caged in an urban stream. *Journal of Neotropical Ichthyology* 5(3):327-336.
- Cambell, Neil A., Reece, Jane B., Mitchell L. G., 2012 [Biology]. Erlangga, Jakarta, 576 p. [In Indonesian].
- Carrion R. L., Alvarellos S. S., Guzman J. M., Martin del Rio M. P., Soengas J. L., Manceraa J. M., 2005 Growth performance of gilthead sea bream conditions: implication for osmoregulation and energy metabolism. *Aquaculture* 250:849-861.
- Dewi, Masithoh, 2013 An effective biofilter type for eel culture in recirculation system. *Elsevier Aquaculture Engineering* 13(2):71-82.
- Febrianti H., Komsanah S., Catur A. P., 2016 [Effect of differences in fatty acid sources on fish meal on the growth of Silver pompano (*Trachinotus blochii*, Lacepede)]. *Journal of Aquawarman* 2(1):24-33. [In Indonesian].
- Fujaya Y., 2004 [Fish physiology]. Jakarta, Rineka Cipta, 179 p. [In Indonesian].
- Hastuti S., Mokoginta I., Dana D., Sutardi T., 2004 [Resistance to stress and immune responses of gouramy (*Osphronemus gouramy*, Lac.) with fish meal containing chromium-yeast]. *Journal Ilmu-Ilmu Perairan Dan Perikanan Indonesia* 11(1):15-21. [In Indonesian].
- Lantu S., 2010 Osmoregulation in aquatic animals. *Jurnal Perikanan dan Kelautan* 6(1):46-50.
- Mc Gavin M. D, Zachary J. F., 2007 Pathologic basic of veterinary disease. Mosby Incorporation, USA, 1476 p.
- Mo Y., 2017 [Looking at the potential of silver pompano]. [www.isw.co.id](http://www.isw.co.id). [In Indonesian].
- Nelvia L., Elfrida, Basri Y., 2015 [Addition of bioball to cultured media filters for survival and growth of carp (*Carassius auratus*)]. [ejurnal.bunghatta.ac.id](http://ejurnal.bunghatta.ac.id). 7(1):1-12. [In Indonesian].
- Nurhidayat K., Nirmala, Djokosetyanto D., 2012 [The effectiveness of biofilter media performance in the recirculation system on water quality for the growth and survival rate of red rainbow (*Glossolepis incisus*, Weber)]. *Journal of Research Aquaculture* 7:89-96. [In Indonesian].
- Pamungkas W., 2012 [Osmoregulation activities, growth responses, and energetic cost in fish reared in a saline environment]. *Media Akuakultur* 7(1):44-51. [In Indonesian].
- Prahadi Y. Y., 2015 [Silver pompano production targeted to grow 31.5%]. *Swa Bisnis*. [In Indonesian].
- Putri A. K., Anggoro S., Djuwito, 2014 Osmotic performance rate and development of silver pompano seeds biomass (*Trachinotus blochii*) cultivated on media with different salinity. *Management of Aquatic Resources* 4(1):159-168.
- Ramsay J. M., Feist G. W., Varga Z. M., Westerfield M., Kent M. L., Schreek C. B., 2006 Whole-body cortisol is an indicator of crowding stress in adult zebrafish, *Danio rerio*. *Aquaculture* 258:565-574.
- Raškoviæ B. S., Stankovi M. B., Markovi Z. Z., Poleksi V. D., 2011 Histological methods in the assessment of different feed effects on liver and intestine of fish. *Journal of Agricultural Sciences* 56(1):87-100.
- Retnani H. T., Abdulgani N., 2013 [Effect of salinity on protein content and growth of Silver pompano (*Trachinotus blochii*)]. *Jurnal Sains Dan Seni Pomits* 2(2):2337-3520. [In Indonesian].
- Rigal F., Thibaud C., Catherine L. N., Guy C., Jean-Antoine T., Fabien A., Patrick B., 2008 Osmoregulation as a potential factor for the differential distribution of two cryptic gobiid species, *Pomatoschistus microps* and *P. marmoratus* in French Mediterranean lagoons. *Scientia Marina* 72(3):469-476.
- Scabra A. R., 2018 [Response of eel fish (*Anguilla bicolor bicolor*) in media with different salinity]. *Jurnal Perikanan* 9(2):180-187. [In Indonesian].
- Setiyoningsih P. R., 2014 [The response of milkfish (*Chanos chanos*) due to changes in salinity with the addition of calcium chloride (CaCl<sub>2</sub>) at different durations]. *Jurnal Penelitian UNISLA* 5(2):6-17. [In Indonesian].

- Siikavuopio S. I., Saether B. S., 2006 Effects of chronic nitrite exposure on growth in juvenile atlantic cod *Gadus morhua*. *Aquaculture* 255:351-356.
- Sitta A., Hermawan T., 2011 [Addition of vitamins and enrichment to live feed to overcome abnormalities of silver pompano seeds (*Trachinotus blochii*, Lacepede)]. Batam Marine Cultivation Center, Directorate of Aquaculture, Ministry of Maritime Affairs and Fisheries, 16 p. [In Indonesian].
- Supriyono E., Supendi A., Nirmala K., 2007 [Utilization of zeolite and activated carbon in the *Corydoras aenus* fish packing]. *Jurnal Akuakultur Indonesia* 6:135-145. [In Indonesian].
- Syakirin M. B., Anggoro S., Prayitno S. B., Saputra S. W., 2018 Effects of the salinity media on the osmotic work level, feed utilization efficiency and the growth of "cantang" hybrid grouper *Epinephelus fuscoguttatus* x *E. lanceolatus*. *AAFL Bioflux* 11(4):1274-1279.
- Takashima F., Hibiya T., 1995 An atlas of fish histology (normal and pathological features). Kodansha, Tokyo, 195 p.
- Thophon S., Kruatrachue M., Upatham E. S., Pokethitiyook P., Sahaphong S., Jaritkhuan S., 2003 Histopathological alteration of white seabass, *Lates calcarifer*, in acute and subchronic cadmium exposure. *Journal of Environmental Pollution* 121:307-320.
- Wahyuni S., Windarti, Putra R. M., 2017 Comparative study on histological structure of gill and kidney of snakehead fish (*Channa striata*, BLOCH 1793) from the Kulim and Sibam Rivers, Riau Province. *Jurnal Online Mahasiswa* 4(2):1-14. [In Indonesian].
- Wulandari A. R., 2006 [The role of salinity in the survival and growth of freshwater silver pompano seed (*Colossoma macropomum*)]. PhD thesis, Fisheries Faculty and Marine Science, Bogor Agricultural Institute, Bogor, Indonesia, 43 p.
- \*\*\* National Standard Indonesia, 2013 [Silver pompano (*Trachinotus blochii*, Lacepede) – Part 4: Seed production]. Jakarta, Indonesia. [In Indonesian].

Received: 08 May 2020. Accepted: 11 June 2020. Published online: 18 June 2020.

Authors:

Mulyadi, Riau University, Faculty of Fisheries and Marine Science, Department of Aquaculture, Jl. H.R. Subantas KM 12.5, 28293 Pekanbaru, Indonesia, e-mail: mulyadibrian26@yahoo.com

Usman Muhammad Tang, Riau University, Faculty of Fisheries and Marine Science, Department of Aquaculture, Jl.H.R. Subantas KM 12.5, 28293 Pekanbaru, Indonesia, e-mail: usman\_mt@yahoo.co.id

Bintal Amin, Riau University, Faculty of Fisheries and Marine Science, Department of Marine Science, Jl.

H.R. Subantas KM 12.5, 28293 Pekanbaru, Indonesia, e-mail: bintalamin@gmail.com

Sukendi, Riau University, Faculty of Fisheries and Marine Science, Department of Aquaculture, Jl.

H.R. Subantas KM 12.5, 28293 Pekanbaru, Indonesia, e-mail: p.sukendims@yahoo.com

Niken Ayu Pamukas, Riau University, Faculty of Fisheries and Marine Science, Department of Aquaculture, Jl.

H.R. Subantas KM 12.5, 28293 Pekanbaru, Indonesia, e-mail: nikenayupamukas@gmail.com

Windarti, Riau University, Faculty of Fisheries and Marine Science, Department of Aquatic Resources

Management, Jl. H.R. Subantas KM 12.5, 28293 Pekanbaru, Indonesia, e-mail: windarti.unri@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Mulyadi, Tang U. M., Amin B., Sukendi, Pamukas N. A., Windarti, 2020 Osmotic performance rate, stress response, and growth performance of silver pompano (*Trachinotus blochii*) reared in different salinities using recirculating culture system. *AAFL Bioflux* 13(3):1546-1556.